# With ProAir® HFA, the savings are BIG

## 15 DAYS until CFC albuterol inhalers banned

### **LINKS**

- www.Switch2HFA.com
- www.ProAirHFA.com
- FDA Public Health Advisory on switching patients now
- Web presentation
  "For a Smooth Switch and Fast-Acting Relief"
- Save your qualified patients up to \$100 off ProAir HFA\*
- Prescribing Information

\*Up to \$20 off their next 5 prescriptions

## The ProAir HFA starter kit provides BIG savings for your patients over Proventil® HFA...

	ProAir HFA	Proventil HFA
Discount Card - Number of Uses	5x (up to \$20)¹	2x (up to \$15) <sup>2</sup>
Voucher for Free Inhaler	YES (only in lieu of sample)	YES (included in starter kit- value of \$35.17) <sup>3</sup>
Total Savings for Patient	\$100	\$65.17

## ...and ProAir HFA is readily accessible on 9 out of 10 commercial managed care plans.<sup>4</sup>

	ProAir HFA	Proventil HFA
Most Preferred Managed Care Access <sup>4</sup>	YES	NO



If you have any questions regarding the Teva Assistance Program™ please call us toll-free at (877) 254-1039.

Learn more about how ProAir HFA supports your switch efforts. Watch "For a Smooth Switch and Fast-Acting Relief – Rely on the #1 Albuterol Brand – ProAir HFA," a Web presentation by a leading pulmonary expert.

### Dispense the #1 albuterol inhaler,<sup>5</sup> ProAir HFA – Now FDA-approved for patients as young as 4

ProAir® HFA is indicated in patients 4 years of age and older for the treatment or prevention of bronchospasm with reversible obstructive airway disease and for the prevention of exercise-induced bronchospasm.

#### Important Safety Information

- Inhaled albuterol sulfate can produce paradoxical bronchospasm that may be life-threatening. It should be recognized that paradoxical bronchospasm, when associated with inhaled formulations, frequently occurs with the first use of a new canister.
- Fatalities have been reported in association with excessive use of inhaled sympathomimetic drugs in patients with asthma.
- ProAir® HFA, as with all sympathomimetic amines, should be used with caution in patients with cardiovascular disorders (especially coronary insufficiency, cardiac arrhythmias, and hypertension), convulsive disorders, hyperthyroidism, and diabetes.
- Potential drug interactions can occur with beta-blockers, diuretics, digoxin, or monoamine oxidase inhibitors, and tricyclic antidepressants.
- Do not exceed the recommended dose.
- Adverse events, which occurred at an incidence rate of at least 3% with ProAir® HFA, include headache, tachycardia, pain, dizziness, pharyngitis, and rhinitis.

REFERENCES: 1. ProAir HFA Discount Card. Available at: https://www.switch2hfa.com/secure\_forms/hcp\_signup.aspx. Accessed September 12, 2008. 2. Proventil Transition Kit. Available at: https://www.proventilhfa.com/phfa/application?names pace=transkit&event=transkit&web\_program\_id=00000000. Accessed September 12, 2008. 3. Analysource June 2008. 4. Business One Technologies. October 1, 2008. 5. IMS Health National Prescription Audit, TRx Data, September 2008.





