

All the benefits of SUPRAX® plus a dosage form for every need

Once-a-day
Suprax®
cefixime
Tablets
400 mg

Once-a-day
Suprax®
cefixime
Oral suspension
200 mg/5 mL

Once-a-day
Suprax®
cefixime
Oral suspension
100 mg/5 mL

Stock up on SUPRAX® now!

Product	Bottle size	NDC#	Ordering Number		
			McKesson	Cardinal	ABC
SUPRAX® Tablets 400 mg	50 tablets	27437-201-08	212-1010	4058897	411-241
SUPRAX® 200 mg/5 mL	50 mL	27437-206-03	162-4972	3970225	659-065
SUPRAX® 200 mg/5 mL	75 mL	27437-206-02	162-5995	3970191	659-072
SUPRAX® 100 mg/5 mL	50 mL	68180-202-03	121-2620	3565728	980-452
SUPRAX® 100 mg/5 mL	100 mL	68180-202-01	121-3107	3565744	980-464



SUPRAX® is indicated for the treatment of uncomplicated urinary tract infections caused by *E. coli* and *P. mirabilis* and uncomplicated gonorrhea (cervical/urethral) caused by *N. gonorrhoeae* (penicillinase- and non-penicillinase-producing strains); otitis media caused by *H. influenzae* (β -lactamase +/- strains), *M. (Branhamella) catarrhalis* (most of which are β -lactamase positive), and *S. pyogenes*; in tonsillitis/pharyngitis caused by *S. pyogenes*.^{*} Otitis media should be treated with suspension only.

Note: Penicillin is the usual drug of choice in the treatment of *S. pyogenes* infections, including the prophylaxis of rheumatic fever. SUPRAX® is generally effective in the eradication of *S. pyogenes* from the nasopharynx; however, data establishing the efficacy of SUPRAX® in the subsequent prevention of rheumatic fever are not available.

Important safety considerations

The most common adverse event related or probably related to SUPRAX® was GI upset. Several patients developed severe diarrhea and/or documented pseudomembranous colitis, and a few required hospitalization. SUPRAX® is contraindicated in patients with known allergy to the cephalosporin group of antibiotics.

WARNINGS: BEFORE THERAPY WITH SUPRAX® IS INSTITUTED, CAREFUL INQUIRY SHOULD BE MADE TO DETERMINE WHETHER THE PATIENT HAS HAD PREVIOUS HYPERSENSITIVITY REACTIONS TO CEPHALOSPORINS, PENICILLINS, OR OTHER DRUGS. IF THIS PRODUCT IS TO BE GIVEN TO PENICILLIN-SENSITIVE PATIENTS, CAUTION SHOULD BE EXERCISED BECAUSE CROSS HYPERSENSITIVITY AMONG β -LACTAM ANTIBIOTICS HAS BEEN CLEARLY DOCUMENTED AND MAY OCCUR IN UP TO 10% OF PATIENTS WITH A HISTORY OF PENICILLIN ALLERGY. IF AN ALLERGIC REACTION TO SUPRAX® OCCURS, DISCONTINUE THE DRUG. SERIOUS ACUTE HYPERSENSITIVITY REACTIONS MAY REQUIRE TREATMENT WITH EPINEPHRINE AND OTHER EMERGENCY MEASURES, INCLUDING OXYGEN, INTRAVENOUS FLUIDS, INTRAVENOUS ANTIHISTAMINES, CORTICOSTEROIDS, PRESSOR AMINES AND AIRWAY MANAGEMENT, AS CLINICALLY INDICATED.

Please [click](#) here for full prescribing information.

 Lupin Pharmaceuticals, Inc.,
Harborplace Tower,
111 South Calvert St., 21st Floor,
Baltimore, MD 21202
A Commitment to Caring for Kids

SUPRAX is a registered trademark.
©2009 Lupin Pharmaceuticals, Inc. December 2009