

YAZ HELPS YOU SAY GOODBYE.

YAZ,® like all birth control pills, is 99% effective, but it's the first and only birth control proven to treat the emotional and physical premenstrual symptoms severe enough to impact your life, a condition doctors call PMDD.

Symptoms could include:

- Irritability
- Moodiness
- Feeling anxious
- Bloating
- Increased appetite
- Fatigue
- Headaches
- Muscle aches

YAZ can also help keep your skin clear. In recent studies, 9 out of 10 women saw improvement in their moderate acne.

YAZ is for women who choose the Pill for contraception. It should be taken as directed, and is available by prescription only.

Who shouldn't take YAZ?

YAZ contains a different kind of hormone that may increase potassium. Therefore, you should not take YAZ if you have kidney, liver, or adrenal disease because this could cause serious heart and health problems. Tell your doctor if you are on daily long-term treatment for a chronic condition such as cardiovascular disease or chronic inflammatory disease. Women who take certain drugs (see below) should have their potassium levels checked in the first month of taking YAZ. **You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.**

Only YAZ goes beyond birth control. Talk to your healthcare professional today to find out if YAZ can help you say goodbye.

YAZ® (drospirenone & ethinyl estradiol)

BEYOND BIRTH CONTROL™

www.YAZ-us.com

1-866-YAZ-PILL

IMPORTANT SAFETY INFORMATION ABOUT YAZ:

What are the risks involved with taking any oral contraceptive (OC)? OCs can be associated with increased risks of several serious side effects. OCs do not protect against HIV infection or other STDs. Women, especially those 35 and over, are strongly advised not to smoke because it increases the risk of serious cardiovascular side effects including blood clots, stroke, and heart attack. What cardiovascular or chronic inflammatory drugs may increase potassium? NSAIDs—ibuprofen (Motrin® Advil®), naproxen (Naprosyn® Aleve® and others) when taken long-term and daily for arthritis or other diseases or conditions, Potassium-sparing diuretics (spironolactone and others), Potassium supplementation, ACE inhibitors (Capoten® Vasotec® Zestril® and others), Angiotensin-II receptor antagonists (Cozaar® Diovan® Avapro® and others), Heparin, and Aldosterone antagonists. See important patient information on reverse side.